



ISSN 0975-413X
CODEN (USA): PCHHAX

Der Pharma Chemica, 2018, 10(2): 36-41
(<http://www.derpharmachemica.com/archive.html>)

Nematicidal Activity of Four Medicinal Plants Extracts against *Meloidogyne* Spp.

Senhaji B^{1*}, Mziouid A¹, Chebli B¹, Mayad EH^{1,2,3}, Ferji Z³

¹Biotechnology and Environmental Engineering Team, National School of Applied Sciences, PO Box: 1136/S, Agadir, Morocco

²Department of Biology, Faculty of Sciences, Ibn-Zohr University, PB. 8106-Cité Dakhla Agadir, Morocco

³Plant Protection Department/Nematology, Horticultural Complex of Agadir, Hassan II Institute of Agriculture & Veterinary Medicine-Agadir, Morocco

ABSTRACT

The aim of this study is to evaluate the effect of organics and aqueous extracts of some aromatic and medicinal plants harvested in Agadir region (Morocco), against second stage juveniles of *Meloidogyne* spp. The four plants used are: *Asteriscus imbricatus*, *Lavendula dentata*, *Pulicaria mauritanica* and *Globularia alypum*. The best results were obtained with *A. imbricatus* extracts: petroleum ether and chloroform extracts induced respectively 89, 31% and 92, 71% of mortality at 2000 ppm, while the aqueous induced 78, 76% of mortality at 20000 ppm. For this reason these extracts have been tested against *Meloidogyne* spp in vivo. The results of in vivo test showed that both nonpolar extracts petroleum ether and chloroform of *A. imbricatus* have an important nematicidal activity at 6000 ppm similar to the effect of chemical nematicid, while the aqueous extract was ineffective. Then the organic extracts of this plant were subjected to a phytochemical screening to get an overview on their metabolites secondary composition. The results showed that the effective extracts in vitro and in vivo are distinguished by the presence of terpenes. The activity demonstrated by *A. imbricatus* was good enough to propose this plant for further studies at greenhouse and field stages to determine its efficacy in soil.

Keywords: *In vitro*, *In vivo*, *Meloidogyne* spp, *Asteriscus imbricatus*, *Pulicaria mauritanica*, *Lavendula dentate*, *Globularia alypum*

INTRODUCTION

Currently the plant pests cause significant losses in crops. These parasites cause yield losses of up to 45% [1] or even a total loss of production. *Meloidogyne* is a phytophagous pest which cause the onset of deformation in the roots of infected plants called galls. *Meloidogyne* parasitize more than 5500 species of plants [2], they are widely spread over the globe and are a dangerous group for the damage it causes [3,4]. Yield losses can reach 65% depending on the level of infestation [5]. The control methods used against these pests are pesticides synthesis, especially methyl bromide remains the most effective remedy for both disinfect the soil nematode that not only to eliminate any kind of terrestrial pathogens and weeds.

Methyl bromide and other nematicidal products are actually withdrawn from the market because of their adverse effects on human health and the environment. Therefore, the search for alternative methods of control, cleaner and cheaper, proves mandatory [6]. The use of plant extracts is one of the solutions that can help reduce the problem of nematodes [6,7]. The Agadir region contains a variety of species having toxic effect against root-knot nematodes. However, although some have been the subject of several studies in the laboratory of Nematology in Hassan II Agriculture and veterinary Institute, Agadir Morocco, but a large number of species have never been evaluated for their toxic effect against root-knot nematodes *Meloidogyne*. It is in this context that fits this study which aims to evaluate the effectiveness of organic and aqueous extracts of four aromatic and medicinal plants against root-knot nematodes *Meloidogyne* spp *in vitro* and *in vivo*, and characterize phytochemically the most effective extracts.

MATERIALS AND METHODS

Collection and preparation of plant

The aerial part of plants samples were collected randomly in April 2012 in three different areas near Agadir (Cape Ghir, Imozzer idaoutanane, Tmanar). The spices of all samples were identified and were deposited in the herbarium in the Mechanics and Process Laboratory, Energy and Environment in National School of Applied Sciences Agadir, Morocco.

The botanical name family, the collecting zone and GPS coordinates are summarized in Table 1. Plant samples were cleaned, and air dried in the

shade then grounded to a fine powder using an electric laboratory grinding mill and stored in the dark at 4°C until use [8].

Table 1: The botanical name family, parts used and the collecting zone of plant samples

Botanical name	Family	The collecting areas	GPS coordinates
<i>Asteriscus imbricatus</i>	<i>Asteraceae</i>	Cape Ghir, 40 km northwest of the city of Agadir	30° 37' 48" N 9° 54' 00" W
<i>Pulicaria mauritanica</i>	<i>Asteraceae</i>	Imozzer idaoutanane, 60 km northwest of the city of Agadir	30° 40' 23" N -9° 28' 48" W
<i>Lavendula dentata</i>	<i>Lamiaceae</i>	Tmanar, 103 km northeast of the city of Agadir	31°00'0 00"N - 9°39'59 99" W
<i>Globularia alypum</i>	<i>globulariaceae</i>		

Preparation of plant extracts

Extraction using organic solvents

We conducted a hot extraction using Soxhlet apparatus, using a cellulose cartridge we filled with plant material powder (Aerial part). Solvents we used are in order of increasing polarity: petroleum ether, the Chloroform, ethyl acetate and methanol [9]. The extraction is complete when the extraction solvent becomes increasingly clear that is to say, without a significant proportion of solute [10]. The final extracts were obtained after concentration and removal of the solvent by rotary evaporation.

Aqueous extraction

The aqueous extraction was performed using 20 g of the powdered herb to 50 ml of distilled water after three hours of maceration using magnetic stirring plaque. The mixture was filtered through filter paper Whatman paper number 4. The remaining cake is recovered and the operation is repeated five times to make five exhaustion. Extracts obtained were stored at 4°C and protected from light until use [11].

In vitro test

Extracting eggs of *Meloidogyne* spp

The roots from a tomato field infested with *Meloidogyne* spp typical galls are rinsed with water, cut into small pieces. Then we proceeded to mix for 2 min in a solution of bleach to 1%. The mixture is filtered through a series of sieves (250 mesh size ranging from microns to 5 µm), and the filtrate containing the nematode eggs are collected on a coarse sieve 2 lined paper handkerchiefs cellulose deposited in a receptacle, the second stage juveniles (J2) of migrating through the cellulose is recovered after 48 h.

Nematicidal assay

Approximately 100 larvae (J2) of *Meloidogyne* spp were placed in an aqueous suspension of 9 ml of H₂O per Petri dish and then added one ml of the extract tested.

The organics extracts were tested against *Meloidogyne* spp, at 20, 200 and 2000 ppm. While the aqueous extract was tested at 20, 200, 2000 and 20000 ppm. Each treatment is carried out in three replicates. The boxes are arranged in a completely randomized device at room temperature (25 ± 2) for 72 h. Distilled water was used as a negative control. During *in vitro* incubation, juvenile (J2) of *Meloidogyne* react differently to plant extracts studied depending on the concentration and the incubation period used (24 h, 48 h and 72 h). Immobile J2 larvae were identified under an optical microscope at a magnification of 40 x. Dead larvae (J2) are identified based on the appearance of needle still they present. A correction is made with respect to the control treatment that distilled water is according to the formula [12].

$$\text{Tim (\%)} = (\text{Tr} - \text{Tm}) / (\text{Tm} - 100) \times 100$$

With, Tr=Percentage of J2 motionless in the extract, Tm=Percentage of J2 immobile in distilled water.

In vivo test

To confirm the results obtained *in vitro*, petroleum ether chloroform and aqueous extracts of *A. imbricatus* plant were selected for *in vivo* test. The experiment was conducted on the culture of tomato at the experimental greenhouse at of Horticultural Complex Agadir, Morocco. The trial lasted from 27/02/2015 (the date of the application of treatment) to 27/04/2015 (date grubbing plants). Plants aged two weeks were transplanted in black plastic pots filled with pretreated substrate by various treatments. The transplantation took place one week after the application of treatments. The transplantation substrate includes 1/3 of peat and 2/3 of a soil naturally infested with *Meloidogyne* spp. We applied eight treatments which are listed in Table 2.

Table 2: The treatments used in vivo

Treatment number	Type of treatment	Concentration(ppm)
T1	Petroleum ether extract of <i>Asteriscus imbricatus</i>	6000
T2	Petroleum ether extract of <i>Asteriscus imbricatus</i>	4000
T3	Chloroform extract of <i>Asteriscus imbricatus</i>	6000
T4	Chloroform extract of <i>Asteriscus imbricatus</i>	4000
T5	Aqueous extract of <i>Asteriscus imbricatus</i>	40000
T6	Aqueous extract of <i>Asteriscus imbricatus</i>	30000
T7	A chemical nematicid: vydate.	5 l/hectare

T8	Positive control consisting of naturally soil infested with <i>Meloidogyne</i> spp but untreated	-
----	--	---

Each treatment was replicated eight times, and pots were arranged in a completely randomized design in a greenhouse.

Phytochemical screening

The extracts which gave the best results *in vitro* and *in vivo*, were selected for phytochemical screening. Each dry residue is dissolved in methanol to prepare a solution of 1 mg/ml. This solution is used to realize all the phytochemical tests.

Characterization of flavonoids

The extracts were subjected to Thin Layer Chromatography (TLC) examination using commercial silica plates for the determination of the secondary metabolites. We tried different solvent systems to achieve better migration and components separation of the studied extracts. For each sample and analysis the same volume 20 μ l was used.

Concerning flavonoids, the best separation of the constituents of studied extracts was obtained using the solvent system toluene / acetic acid / water (125:72:3) [13]. The revelation of different fluorescences was carried out under UV light at 365 nm after spraying with the NEU reagent (2 amino ethyl diphenylboric). The frontal report (Rf) was calculated as: The distance line-compound deposition on the line distance of solvent.

Characterization of terpenes and alkaloids

For terpenes, the benzene was used as a migration solvent. After migration, the silica gel plate was sprayed with antimony chloride, and then placed in an oven at 110°C for 10 min. Any fluorescence detected, under UV at 365 nm, after this treatment proves that the tested material contains terpenes (Randerath 1971). For alkaloids we used the solvent system consisting of (AcEt/MeOH/NH₄OH) (9/1/1). The detection was carried out by spraying the Dragendorff reagent. The presence of alkaloids was shown by a bright orange color on the chromatogram [14].

Test for tannins

A few drops of 0.1% ferric chloride was added to 5 each organic extract and observed for brownish green or a blue-black coloration [15].

Statistical analysis

The data were statistically analysed using SPSS 16.0 software.

We performed an analysis of variance (ANOVA). Student-Newman-Keuls tests were used to segregate treatments which were significantly different at $P < 0.05$.

RESULTS AND DISCUSSION

Results of *in vitro* tests

Nematicidal activity of organics extract against *Meloidogyne* spp

The results presented in Table 3 show the nematicidal activity of organic extracts of four plant species belonging to three botanical families, against *Meloidogyne* spp, after 72 h evaluated at 20, 200 and 2000 ppm. Among the four species tested, three had some nematicidal activity. *cc*, *A. imbricatus* and *L. dentata*. While the fourth plant *G. alypum* showed no effect against *Meloidogyne* spp. Organics extracts of *A. imbricatus* showed the highest nematicidal activity (mortality rate ranging from 2,41 to 92,71%) followed by *P. mauritanica* plant who showed moderate activity (mortality rate ranging from 2,21 to 37,5%) and finally *L. dentata* who showed low activity (mortality rate ranging from 0,08 to 2,32%).

The statistical analysis for each plant extract according to the concentration (results obtained in each column) showed a significant difference between the concentrations tested. Indeed the mortality rate of *Meloidogyne* spp increases with increasing concentration. In addition the interesting results were observed at 2000 ppm.

Table 3: *In vitro* effects of organic plants extracts on mortality rate (%) of *Meloidogyne* spp, after 72 h evaluated at 20,200 and 2000 ppm

Concentration (ppm)	<i>Asteriscus imbricatus</i>				<i>Pulicaria mauritanica</i>				<i>Lavendula dentata</i>				<i>Globularia alypum</i>			
	PE	CHL	ACT	MT	PE	CHL	ACT	MT	PE	CHL	ACT	MT	PE	CHL	ACT	MT
20	2,41 ^c	2,95 ^c	-3,78 ^b	-3,73 ^b	4,19 ^b	2,21 ^b	0	0	0	0	0	0	0	0	0	0
200	45,2 ^b	45,88 ^b	4,71 ^b	5,57 ^{ab}	6,71 ^b	11,43 ^b	3,64 ^b	0	0,08	0	0	0	0	0	0	0
2000	89,31 ^a	92,71 ^a	36,24 ^a	17,42 ^a	37,05 ^a	36,35 ^a	19,47 ^a	13,8 ^a	1,31	2,32	0,66	0	0	0	0	0

(PE): Petroleum ether; (CHL): Chloroform; (ACT): Ethyl acetate; (MT):Methanol; C: Concentration of organic extracts.

Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at $P < 0.05$ according to Newmanand Keuls test. In a second statistical analysis for each plant we evaluated the influence of the type of extract on the mortality of *Meloidogyne* spp at the same concentration of 2000 ppm (The results are represented in each column of Table 4).

A significant difference was observed between the extracts tested. The Newmanand Keuls test has shown the presence of at least two homogeneous groups for both *Asteraceae* plants: *A. imbricatus* and *P. mauritanica*. The first one is constituted by no polar extracts petroleum ether and chloroform. While the others groups are constituted by ethyl acetate and methanol extracts, which means that the mortality rate of *Meloidogyne* spp increased with decreasing polarity.

Table 4: A mortality rate (%) of *Meloidogyne* spp at 2000 ppm depending on the type of organic extracts

Concentration extracts (ppm)	<i>Asteriscus imbricatus</i>	<i>Pulicaria mauritanica</i>	<i>Lavendula dentata</i>	<i>Globularia alypum</i>
------------------------------	------------------------------	------------------------------	--------------------------	--------------------------

Petroleum ether	89, 31 ^a	37,05 ^a	1,31 ^a	0
Chloroform	92, 71 ^a	36,35 ^a	2,32 ^a	0
Ethyl acetate	36, 24 ^b	19,47 ^b	0	0
Methanol	1,742 ^c	13,8 ^b	0	0

Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at $P < 0.05$ according to Newman and Keuls test.

Effect of aqueous extracts against *Meloidogyne* spp

Table 5 shows the results of the effect of aqueous extracts of four plants studied on the mortality percentage of *Meloidogyne* spp after 72 hours, evaluated at different concentrations (20000, 2000, 200 and 20 ppm). Exposure of J2 larvae of *Meloidogyne* spp during 72 h to the aqueous extracts of these four plants at 20000 ppm, showed a different nematicidal effect: the higher nematicidal activity was detected with the aqueous extracts of *A. imbricatus* 78.76% of mortality, followed by the aqueous extract of *P. mauritanica* 9.5% of mortality while the two plants *Lavandula dentata* and *G. alypum* showed no nematicidal effect (Table 5).

Table 5: In vitro effects of aqueous extracts on mortality rate (%) of *Meloidogyne* spp

Concentration (ppm)	<i>Asteriscus imbricatus</i>	<i>Pulicaria mauritanica</i>	<i>Lavendula dentata</i>	<i>Globularia alypum</i>
20	0 ^c	0 ^b	0	0
200	2,25 ^c	1,11 ^a	0	0
2000	14,99 ^b	1,94 ^a	0	0
20000	78,76 ^a	9,5 ^a	0	0

Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at $P < 0.05$ according to Newman and Keuls test.

Based on the results of *in vitro* tests, petroleum ether, chloroform and aqueous extracts of *A. imbricatus* were selected to testing their effect on the damage of tomato plants which is manifested by the appearance of galls on roots.

Results of *in vivo* test

The effectiveness of various treatments against *Meloidogyne* spp *in vivo* was evaluated based on the index of gall. At the end of the test plants were uprooted and roots were carefully washed with tap water. The number of galls on roots of each treatment was counted. Then the gall index was determined based on a 0-5 scale [4]. A significant difference was observed between the treatments tested. The Newman and Keuls test has shown the presence of two homogeneous groups. The first one is constituted by petroleum ether and chloroform extracts at 6000 ppm in the addition of chemical nematicid. The second group contains the positive control and the other treatments (Figure 1).




Figure 1: Effect of petroleum ether, chloroform and aqueous extracts of *Asteriscus imbricatus* on the index gall

Results of phytochemical screening

Flavonoids characterization using TLC

The results of flavonoids characterization of *A. imbricatus* organic extracts are represented in the Figure 2 (Picture 1). The analysis of TLC under UV light after revelation shows: Firstly the fluorescence appears in the four organic extracts, showing that they all contain flavonoids but in various proportions. The ethyl acetate extract has more bands with intense fluorescence compared to other extracts. Secondly this fluorescence is intensified when the polarity of the solvent increases, from petroleum ether chloroform and ethyl acetate extract, but it decrease for the methanol.

Terpenes and alkaloids characterization

For this plant, a blue fluorescence is observed in both petroleum ether extract and chloroform extract which means that these extracts contain terpenes while the two other extracts do not contain terpenes Figure 2 (Picture 2). The presence of alkaloids is demonstrated by spraying the chromatogram with Dragendorff reagent. For *A. imbricatus*, we notice the appearance of orange spot in chloroform and ethyl acetate extracts, which indicate that these extracts contain the alkaloids. While the petroleum ether and methanol extracts showed no stain Figure 2 (Picture 3).

Tannins


The presence of tannins was observed at the ethyl acetate and methanol extracts which showed brownish green coloration. The results of the

phytochemical screening are summarized in Table 6.

Table 6: Phytochemical constituents of *Asteriscus imbricatus* organic extracts

	Flavonoids	Terpenes	Alkaloids	Tannins
Petroleum ether	+	+	-	-
Chloroform	+	+	+	-
Ethyl acetate	+	-	+	+
Methanol	+	-	-	+

+: Present - : Absent



1: Flavonoids characterization; 2: Terpenes characterization; 3: alkaloids characterization
E: Petroleum ether; C: Chloroform; A: Ethyl acetate; M: Methanol

Figure 2: Phytochemical characterization of *Asteriscus imbricatus* organic extracts

DISCUSSION

The results of *in vitro* and *in vivo* experiments show that the *A. imbricatus* plant is more effective against *Meloidogyne* spp compared to the other species tested. These results are consistent with several previous researches that has shown the effectiveness of Asteraceae plants to controlling nematodes: *Tagetes patula* for example has long been known as nematicidal plant [16-18]. Nematicidal activity of *Artemisia* genus was reported by some studies [19]. The effectiveness of plants against nematodes is due to their richness in secondary metabolites. Indeed several previous researchers proved nematicidal activity of diverse classes of these compounds: Saponins [20], alkaloids [21,22], phenolics [23], polyacetylenes, fatty acids, terpenoids [24,25], and others. For Asteraceae plants some nematicidal compounds have been isolated. The most studied are Polythienyls, especially α -terthienyl isolated from *Tagetes* species [26-29]. For this reason, the *A. imbricatus* organic extracts of this plant have been phytochemically screened to see what are the compounds contained in these extracts and how they relate to the activity studied. The results of this study showed that flavonoids are present in the four organic extracts of *A. imbricatus*. The terpenes are present only in petroleum ether and chloroform extracts, the alkaloids are present in chloroform and ethyl acetate extract and finally tannins are present in ethyl acetate and methanol extracts. The analysis of all the results revealed a correlation between the phytochemical content of these extracts and the nematicidal activity. Indeed a remarkable correlation appears between the nematicidal power of the extracts and the presence of the terpenes: the two extracts of petroleum ether and chloroform which contain the terpenes gave a strong nematicidal activity, whereas this activity was very low for the two extracts d ethyl acetate and methanol which do not contain. Terpenes have been reported to possess nematicidal properties: β -sitosterol, stigmasterol, daucosterol, carvacrol, thymol, geraniol, camphor and linalool [29], Gossypol, hemigossypol, methoxygossypol, methoxyhemigossypol dimethoxygossypol [25]. Borneol, carvedel, citral, geraniol, and α -terpineol [30].

Our phytochemical study revealed also the presence of flavonoids in petroleum ether and chloroform which have given a better nematicidal effect. These results suggest that flavonoids may contribute to nematicidal effect. In this context several works in the literature have proved the effectiveness of these metabolites against nematodes: Flavonoids are synthesized by leguminous plants in response to infection by nematode [31]. glyceolline inhibits the oxidation, respiration and motility of *Meloidogyne incognita* [32]. González et al. have shown that the (E)-chalcone (trans-1,3-diphenylpropenone) inhibited the eggs hatching of *Globodera pallida* et *Globodera rostochiensis*. isoflavonoïds like quercétin are nématotoxic to *Meloidogyne incognita* [33,34].

CONCLUSION

Nematicidal *In vitro* and *in vivo* tests showed that both petroleum ether extracts and chloroform of *A. imbricatus* plant have an interesting nematicidal power. The phytochemical screening, realized by chromatographic characterization, showed the richness of these two extracts into different secondary metabolites essentially terpenes and flavonoids. Who are probably responsible for this activity. To confirm this hypothesis, further studies are needed to complete our results, especially the evaluation of the isolated and combined effect of these metabolites against *Meloidogyne* spp *in vitro* and *in vivo*. It would also be interesting to test these extracts at greenhouse and field stages to evaluate their effectiveness against nematodes in soil, and based on the results obtained, the plant could be recommended for use in agriculture in the future. Finally the results of this work, encourage the investigations of aromatic and medicinal plants from the Souss-Massa region.

REFERENCES

- [1] J. Sasser, *J Nematology.*, **1977**, 9, 1, 26.
- [2] V. C. Blok, J. T. Jones, M. S. Phillips, D. L. Trudgill, *Bio Essays.*, **2008**, 30, 249.
- [3] C. Netscher, Les nématodes parasites des cultures maraîchères au Sénégal, Cahiers ORSTOM Série Biologie., **1970**, 11, 209.
- [4] J. Sasser, M. Kirby, **1979**.
- [5] A. T. Ploeg, M. S. Phillips, *Nematology.*, **2001**, 3, 151.
- [6] Y. Oka, *Nematology.*, **2001**, 3, 2, 159.
- [7] M. Takasugi, Y. Yachida, M. Anetai, T. Masamune, K. Kegasawa, *Chemistry Letters.*, **1975**, 4, 1, 43.
- [8] I. Talibi, L. Askarne, H. Boubaker, E. Boudyach, F. Msanda, B. Saadi, A. Aoumar, *Crop Protection.*, **2012**, 35, 41.
- [9] N. Dohou, K. Yamni, K. Badoc, A. Douira, *Bull. Soc. Pharm Bord.*, **2004**, 143, 31.
- [10] S. Harnett, V. Oosthuizen, M. Van de Venter, *J Ethnopharmacol.*, **2005**, 96, 1, 113.
- [11] O. Senhaji, M. Faid, M. Elyachioui, M. Dehhaoui, *J Medical Mycology.*, **2005**, 15, 4, 220.
- [12] W. Abbott, *J. Econ. Entomol.*, **1925**, 18, 2, 265.
- [13] K. R. Markham, Techniques of flavonoid identification, London: Academic press, **1982**, 31.
- [14] K. Randerath, Chromatographie sur couches minces, Paris, **1971**, Édition Gauthier-Villars, 337.
- [15] G. Ayoola, H. Coker, S. Adesegun, A. Adepoju-Bello, K. Obaweya, E. Ezennia, T. Atangbayila, *Tropical. J. Pharm. Res.*, **2008**, 7, 1019.
- [16] J. Good, N. Minton, C. Jaworski, *Phytopathology.*, **1965**, 55, 9, 1026.
- [17] G. Poinar, P. Johnson, How to neutralize nematodes. Rodale's organic gardening, USA, **1986**.
- [18] L. Reynolds, J. Potter, B. Ball-Coelho, *Agronomy J.*, **2000**, 92, 5, 957.
- [19] E.H. Mayad, Z. Ferji, L.M. Idrissi, B. Chebl, *Reviews in Biol Biotechnol.*, **2006**, 5, 2, 37.
- [20] M. P. Argentieri, T. D'Addabbo, A. Tava, A. Agostinelli, M. Jurzysta, P. Avato, *Europ. J. Plant. Pathology.*, **2008**, 120, 2, 189.
- [21] B. Zhao, *J. Chem. Ecol.*, **1999**, 25, 10, 2205.
- [22] T. Thoden, M. Boppré, *Nematology.*, **2010**, 12, 1, 1.
- [23] R. Mahajan, D. Kaur, K. Bajaj, *Nematologia Mediterranea.* **1992.**, 20, 2, 217.
- [24] J. Veech, *J Nematology.*, **1977**, 11, 3, 240.
- [25] N. Ntalli, F. Ferrari, I. Giannakou, U. Menkissoglu Spiroudi, *Pest Management Science.*, **2011**, 67, 3, 341.
- [26] C. Wat, S. Prasad, E. Graham, S. Partington, T. Arnason, G. Towers, J. Lam, *Biochemical Systematics and Ecology.*, **1981**, 9, 1, 59.
- [27] F. Gommers, J. Bakker, *Diseases of Nematodes.*, **1988**, 1, 3, 22.
- [28] D.J. Chitwood, Annual review of phytopathology., **2002**, 40, 1, 221.
- [29] P. Ohri, S. K. Pannu, *Environmental Res. Development.*, **2009**, 4, 171.
- [30] S. Echeverrigaray, J. Zacaria, R. Beltrão, *Phytopathology.*, **2010**, 100, 199.
- [31] P. Hutangura, U. Mathesius, M. G. Jones, B. Rolfe, *Funct. Plant Biol.*, **1999**, 26, 221.
- [32] D. Kaplan, N. Keen, I. Thomason, *Physiological Plant Pathol.*, **1980**, 16, 309.
- [33] M. Begum, M. Sivakumar, *Nematol.*, **2004**, 15, 73.
- [34] T. Tsay, S. Wu, Y. Lin, *J. Nematol.*, **2004**, 36, 1, 36.